

MR IMAGING OF PRIMARY MALIGNANT BONE TUMOURS WITH HPE CORRELATION

R Vikash Babu¹, M Rajya Lakshmi²

Received : 29/07/2025
 Received in revised form : 13/09/2025
 Accepted : 02/10/2025

Keywords:

Bone tumours, osteosarcoma, osteoblastoma, Giant cell tumour, Enchondroma, chondrosarcoma, chondroma histopathology, radiographic features, MRI features, CT findings.

Corresponding Author:

Dr. M Rajya Lakshmi,

Email: ranri2k9@gmail.com

DOI: 10.47009/jamp.2025.7.5.107

Source of Support: Nil,

Conflict of Interest: None declared

Int J Acad Med Pharm
 2025; 7 (5); 543-550



¹Assistant Professor, Department of Radio Diagnosis, Konasemainsistute of Medical Sciences, Amalpuram, Dr B R Ambedkarkonasema District, Andhra Pradesh, India.

²Assistant Professor, Department of General Medicine, Konasemainsistute of Medical Sciences, Amalpuram, Dr B R Ambedkarkonasema District, Andhra Pradesh, India.

ABSTRACT

Background: Aims and Objectives: 1. The purpose of this study is to evaluate the role of MRI in cases of primary malignant bone tumours and MRI characteristics of different primary malignant bone tumours. 2. To compare the imaging findings with surgical and gross pathological findings. 3. Staging of tumour on MRI, correlating them with operative and histopathological findings. **Materials and Methods:** This was a prospective study done in Department of Radiodiagnosis of NRIGH comprising of 40 patients 21 male and 19 female who are suspected or proven cases of the malignant bone tumor. Plain radiographs in AP and Lateral views (including the adjacent joint) are taken in all cases. **MRI Protocol:** The primary pulse sequences include T1 and T2 WI using spine echo and gradient echo techniques with TR of 600 msec. and TE 30msec. for T1WI and TR of 2740 msec. and TE of 85 msec. for T2 WI. The MR morphology was correlated with surgical and histopathological features. **Result:** The study “Multiplanar MR Imaging of primary malignant bone tumours with surgical and histopathological correlation” comprised of 40 patients in a two-year period starting from august 2010 to September 2012 the age ranged from 8 years to 71 years (mean 40 yrs). There were 21 males and 19 females. **Conclusion:** MRI in combination with plain radiography is an excellent modality for evaluation of the musculoskeletal pathologies especially differentiating a malignant from a benign lesion. The multiplanar imaging capabilities place a major role in delineation of tumour extent in to the bone and soft tissues with high contrast and resolution with additional information of neurovascular bundle involvement, joint involvement and staging.

INTRODUCTION

A bone tumour,^[1] is a neoplastic growth of tissue in bone. Abnormal growths found in the bone can be either benign (noncancerous) or malignant (cancerous).

Cross sectional imaging had improved the diagnostic accuracy of both benign and malignant bone tumors. The ability to characterize tumors, the differential diagnosis of primary osseous neoplasm remains based on the radiographic appearance. Radiographs provide information regarding the location, margin, matrix mineralization, cortical involvement and adjacent periosteal reaction. But MRI is superior to plain radiography due to its delineation of local extent and local staging of the disease. And also in assessing the intraarticular extension and the presence of intratumoral necrosis and haemorrhage. In assessing the treatment effectiveness and follow-up.

MATERIALS AND METHODS

This was a prospective study done in Department of Radiodiagnosis of KIMS comprising of 40 patients 21 male and 19 female who are suspected or proven cases of the malignant bone tumour. Plain radiographs in AP and Lateral views (including the adjacent joint) are taken in all cases. MRI was performed on a 1.5 Tesla GE system. The primary pulse sequences included T1 and T2W1 using spin echo and gradient echo techniques. The additional sequences used include the fat saturated SE and short Tau Inversion-Recovery (STIR) sequences. MRI was performed on a 1.5 Tesla GE (General electrical) Signa Excite. The primary pulse sequences include T1 and T2 WI using spine echo and gradient echo techniques with TR of 600 msec. and TE 30msec. for T1WI and TR of 2740 msec. and TE of 85 msec. for T2 WI. Additional sequences used include the FS SE and STIR sequences.

The spinal cord is imaged in sagittal, coronal and axial planes. Images are obtained with multi slice technique using slice thickness using a slice thickness of 5mm, interslice gap of 6mm and matrix of size of 512 x 512.

The MR morphology was correlated with surgical and histopathological features.

All the Giant cell tumours were included in the study because it was not possible to predict the malignancy on plain radiography.

The finding to be noted are given in Table 1

1. The transitional zone
2. Intramedullary extent
3. Soft tissue extent
4. Mineralization of matrix
5. Periosteal response
6. Cortical involvement
7. Joint involvement
8. Site of lesion
9. Epiphyseal involvement
Features seen only on MRI
1. Involvement of neurovascular bundle
2. Signal characteristics.

RESULTS

The study “Multiplanar MR Imaging of primary malignant bone tumours with histopathological

correlation” comprised of 40 patients in a two-year period starting from november 2017 to january 2019 the age ranged from 6 years to 71years (mean 40 yrs). There were 21 males and 19 females.

Study Group

Table 1

Osteosarcoma	13	7	6	22-Jun	14
Ewing's sarcoma	8	4	4	29-Sep	19
Chondrosarcoma	6	3	3	28-71	49
Chordoma	3	3	0	34-64	49
GCT	8	2	6	21-50	35
Multiple myeloma	2	2	0	54-65	60

The common clinical symptoms were pain in the region of tumour and swelling lasting for 2-3 months on average. Three cases gave history of antecedent trauma.

All the patients were investigated with plain radiographs and MRI and were confirmed with gross surgical and histopathological findings.

Histopathological Classification

Osteosarcoma	13
Ewing's sarcoma	8
Chondrosarcoma	6
GCT	8
Multiple Myeloma	2
Chordoma	3

Of the 40 cases, thirteen were Osteosarcomas, eight were Ewing's sarcomas, six were Chondrosarcomas, eight were Giant cell tumours, three were

Chordomas, and two were multiple myelomas. Of the eight Giant cell tumours, one case was malignant.

Osteosarcomaewing's Sarcoma

Lower end of Femur	4
Upper end of Tibia	3
Upper end of Femur	2
Upper end of Humerus	3
Thigh	1
Shaft of femur	3
Ilium	2
Femur upper end	1
Humerus upper end	1
Rib	1

GIANT CELL TUMOURCHONDROSARCOMA

Radius lower end	2
Humerus upper end	2
Femur upper end	1
Tibia upper end	1
Tibia lower end	2

Pelvis	3
Femur	2
Humerus upper end	1

CHORDOMA

SACRUM	3
--------	---

MRI FINDINGS IN MALIGNANT BONE TUMOURS




Signal Characteristics

Most of the tumours were hypointense on T1W images and heterogeneously hyperintense on T2W images. In general most tumours had a non specific appearance on MRI except for cartilaginous tumours. The cases of chondrosarcoma were profoundly hyperintense on T2W images because of high water content of cartilaginous elements.

Cortical break


Of the total 40 cases cortical break was detected on MRI in 33 cases. It was absent in 7 cases. Thus 82.5%cases demonstrated cortical break and 17.5% did not show cortical break on MRI. It is best demonstrated on T1W imaging. Thirty one cases were operated and nine cases were not operated due to the presence of distant metastasis. Out of the thirty one cases operated, cortical involvement was seen in 27 cases and was absent in 4cases.

CORTICAL INVOLVEMENT	MRI FINDINGS	SURGICAL FINDINGS
PRESENT	26/31 (84%)	27/31 (87%)
ABSENT	5/31 (16%)	4/31(13%)



Site of the Lesion

Cortical Break



Thus sensitivity, specificity, positive predictive value and negative predictive value of MRI in detecting cortical involvement in our study are 96.2%,100%,100% and 80% respectively.

Marrow involvement


Marrow involvement was seen in all cases on MRI which was confirmed on surgery in thirty one cases. The extent of marrow involvement was best shown by T1W images and STIR coronal or sagittal sequence. It was confirmed by surgical and pathological findings in thirty one cases.

Soft tissue involvement:

Extra osseous soft tissue involvement was seen in 34 cases out of 40 cases. Soft tissue involvement is not seen in 4 cases of GCT, two cases of Ewing's sarcoma. Extraosseoussofttissue involvement was best shown by T2W axial images.

SOFT TISSUE INVOLVEMENT	MRI FINDINGS	SURGICOPATHOLOGICAL FINDINGS
PRESENT	24/31 (77.4%)	25/31 (80.6%)
ABSENT	7/31 (19.3%)	6/31 (16%)

(No. of Cases)



involvement of joint and 67.5% cases did not show joint involvement on MRI.

Surgery could only be performed in 31 of these patients due to presence of metastasis in nine cases.


Joint involvement

Out of total forty cases, 12 cases showed joint involvement on MRI. The joint was uninvolved in 27 cases. Thus 32.5% cases demonstrated

JOINT INVOLVEMENT	MRI FINDINGS	SURGICO PATHOLOGICAL FINDINGS
PRESENT	12/31 (38.7%)	10/31 (32.2%)
ABSENT	19/31 (61.2%)	21/31 (67.7%)

The sensitivity was 100%, specificity 90.4%, positive predictive value 83.3% and negative predictive value 100%.

(No. Of Cases)



Out of a total of forty cases, MRI showed neurovascular bundle involvement in four cases. It was uninvolved in 36 cases. Thus 10% cases demonstrated involvement of neurovascular bundle and 90% cases did not show involvement of neurovascular bundle on MRI. Surgery could be performed in 31 of these patients. Nine patients could not be operated due to the presence of metastasis.


MRI and surgical/pathological correlation out of 35 patients who were operated was as follows

Neurovascular bundle involvement

NEUROVASCULAR INVOLVEMENT	MRI FINDINGS	SURGICAL/PATHOLOGICAL FINDINGS
INVOLVED	4/31 (12.9%)	5/31 (16%)
UNINVOLVED	27/31 (87.1%)	26/31 (84%)

The sensitivity in our study is 100%, specificity 96.2%, positive predictive value 100 % and negative predictive value is 96.2%

Neurovascular Bundle Involvement



DISCUSSION

In our study, we tried to prospectively analyse the accuracy of MRI in relative to conventional radiography. We studied forty cases of primary bone tumours (thirteen Osteosarcomas, eight Ewing's sarcomas, six Chondrosarcomas, eight osteoblastomas, three chordomas and two multiple myelomas) by radiography and MR imaging and analysed the findings.

On plain radiography and MRI, the parameters mentioned in the table 1 were evaluated in all the and staging was done separately.

Transitional zone

Of the thirteen cases of Osteosarcoma, narrow zone was noted in three and the ten cases showed wide zone of transition on plain radiography. All the cases of Ewing's sarcoma and Chondrosarcoma showed wide zone of transition. All the eight cases of giant cell tumour showed narrow zone of transition.

On MRI, all the lesions showed clear demarcation between the involved zone and normal fat in the marrow. The giant cell tumours showed a clear hypointense rim corresponding to the sclerosis on plain radiograph.

Marrow Involvement

In our study marrow involvement was seen in all the forty cases. It was confirmed by surgical and pathological findings in 31 cases. The extent of marrow involvement was best shown by T1W images, STIR coronal or sagittal sequences.

Ella Onikul,^[2] David Parham et al in 1996 studied the accuracy of MR imaging for estimating intraosseous extent of osteosarcoma. They compared how well T1W images and STIR images revealed the extent of longitudinal intraosseous involvement in osteosarcoma.

R Golfieri et al,^[38] in 1990 studied the role of STIR sequence in MRI examination of bone tumours and found that the STIR sequence suppress the high signal from fatty bone marrow giving a clear depiction of tumour extent in its intramedullary component.

Soft tissue component

Soft tissue involvement was seen in 34 out of 40 cases. Two cases of Ewing's sarcoma did not show any obvious soft tissue component. Extra osseous involvement was best shown by T2 weighted axial images.

Orest B Boyko and David A Cory,^[39] in 1986 evaluated 25 patients with Osteogenic sarcoma and Ewing's sarcoma with MRI and found that tumour involvement of the soft tissue is best shown by T2 weighted sequences.

Cortical Break

Cortical break was detected on MRI in 33 cases. It was absent in 7 cases. It is best demonstrated on T1 longitudinal and T1 axial images. Thirty-one cases were operated. Nine cases could not be evaluated due to the presence of metastases. Out of thirty-one cases operated cortical involvement was noted in 26 cases and absent in 5 cases.

The sensitivity, specificity, positive predictive value and negative predictive value of MRI in detecting cortical involvement in our study are 96.2%, 100%, 100% and 80% respectively.

Joint involvement

The presence or absence of joint involvement is particularly important in preoperative evaluation of tumour extent which will subsequently decide the appropriate surgical procedure (intra or extra articular resection). MRI is highly sensitive for detecting joint involvement may lead to overstating

of tumour and result in unnecessarily radical surgical procedures.

(Wolfgang and Schima 1994).^[40]

We evaluated the relationship of the tumour to adjacent joint and involvement of the intrasynovial joint space was presumed where T1W images showed that a contrast enhancing mass extended into the joint space either by disruption of joint capsule or by intraarticular destruction of cortical bone and the articular cartilage, if present. Joint involvement was also presumed in cases of tumour extension into the cruciate ligaments which are intracapsular but extrasynovial. The presence of a joint effusion did not allow differentiation with regard to tumour extension into the joint. Only the absence of a joint effusion had a high predictive value for absence of joint involvement by tumour. Compared with unenhanced T1 weighted images, contrast enhanced T1 weighted images are more useful for assessing joint involvement. When MR imaging is used, the presence of peritumoural inflammatory changes may lead to false positive diagnosis of joint involvement which may be followed by unnecessary radical en bloc resection of joint.^{(Schima 1994).^[30]}

In our study MRI showed joint involvement in 12 cases, it was uninvolved in 19 cases. surgery was performed in 31 cases due to the presence of metastasis in nine cases. The sensitivity was 100%, specificity 90.4%, positive predictive value 83.3% and negative predictive value 100%.

The results in our study are quite similar to Van Trommel et al,^[41] who found sensitivity, specificity, positive predictive value and negative predictive value to be 100%, 70%, 86.4% and 100% respectively.

Neurovascular bundle involvement

Radiologic depiction of soft tissue mass with respect to neurovascular bundle is important in planning surgical approaches for local tumour control. Tumour with deviation of neurovascular bundle may be considered for amputation instead of limb salvage procedure because the ability to obtain adequate surgical margins around the tumour may be compromised.

On MRI involvement of neurovascular bundle was present when tumour is surrounding these structures or containing at least one half the circumference and obliterating the associated fat plane. The relationship of neurovascular bundle to the tumour was best shown on T2 weighted axial images and T1 weighted post contrast axial images. Fat saturated T1 weighted post contrast images are superior to T2 weighted images in defining the proximity of soft tissue tumour mass to the neurovascular bundle. Suzanne, William Kaufman,^[42] in 1997 found that it is easier to evaluate neurovascular bundle proximity to tumour with fat sat T1 post contrast images than with T2W for 64% of cases. In our study MRI showed neurovascular bundle involvement in four cases. It was uninvolved in 36 cases. Surgery could

be performed in 31 cases. Nine patients could not be operated due to the presence of metastasis. One case was detected false positive on MRI. The sensitivity in our study is 100%, specificity 96.2%, positive predictive value 100 % and negative predictive value is 96.2%. The result in our study is comparable to Bloem JL et al,^[43] who found sensitivity, specificity, positive predictive value and negative predictive value to be 100%, 98%, 98% and 100% respectively.

Signal intensity pattern

Despite great value of MR imaging in the staging of the bone lesions, it is of relatively little value in specific histological diagnosis. There are specific diagnoses however that have a relatively characteristic MR appearance. Cohen et al,^[27] observed a distinctive MR appearance in chondroid lesions containing a matrix of hyaline cartilage. The unique pattern consisted of homogenous high signal in a discernible lobular

configuration on T2 weighted spin echo images. This MR appearance in chondroid lesions reflects underlying high ratio of water content to mucopolysaccharide component within hyaline cartilage. MR imaging also allows precise measurement of thickness of cartilage cap of an osteochondroma. It is agreed that the risk of malignant transformation is directly related to thickness of cartilage cap especially when later exceeds 2 cm. In our study, six cases of chondrosarcomas were correctly characterised with MRI. They were profoundly hyperintense on T2 weighted images because of high water content of cartilaginous elements. Rest of the tumours had a non-specific appearance on MRI.

Staging of tumours

Enneking W F, Spanier S S, Goodman M A,^[11] proposed a system for the surgical staging of musculoskeletal sarcomas.

STAGE	GRADE	SITE	METASTASES
IA	G1	T1	M0
IB	G1	T2	M0
IIA	G2	T1	M0
IIB	G2	T2	M0
IIIA	G1-2	T1	M1
IIIB	G1-2	T2	M1

Characteristics:

G- Grade T-Site

G0- benign T1- Intracompartmental

G1-low grade malignancy T2-extracompartmental

G2-high grade malignancy

M- Metastasis

M0-no regional or distant metastasis

M1- regional or distant metastasis present.

The malignant tumours were staged according to the Enneking's Staging of Musculoskeletal Neoplasms.^[11] Nine cases had distant metastasis (Stage III), 21 cases were of Stage IIB and 2 cases were in Stage IIA.

STAGE IIA	STAGE IIB	STAGE III
2	21	9

The Giant Cell Tumours were staged according to Enneking's modified staging system:

Stage 1: asymptomatic and latent Radiographically with benign histologic pattern.

Stage 2: may be symptomatic but there is "active" radiographic appearance (the GCT has expanded the affected bone but is still confined by the overlying cortex.), there is no radiographic evidence of distant metastasis with benign histology.

Stage 3: is symptomatic and radiographs show rapid and aggressive growth (permeative bone destruction, extracortical extension with accompanying soft tissue mass, extension to the subcortical cortex and/or associated periosteal reaction). The histologic pattern is benign and distant metastases are rare but acceptable for this stage.

According this staging system six cases were instage II and two cases were in stage III in our study.

The discrepancy of staging between plain radiography and MRI was mainly due to the cortical invasion and soft tissue swelling evident on MRI alone. According to Murali Sundaram et al when

radiographic depiction of tumour permits assessment of its morphology, matrix and probable histologic nature, MRI ought to be the next examination solely for staging purposes.^[20] Bloem et al also concluded in their study of 56 patients with bone sarcomas that MR imaging was the modality of choice for local staging of primary bone sarcoma.^[49]






Figure 1: 8 yrs./M child c/o swelling in the right shoulder since 3 months A: X-ray Rt shoulder shows Fracture neck of right humerus with large soft tissue opacity noted in the upper end of humerus with calcifications, sclerosis and periosteal reaction which are confirmed by the axials of CT (B) and axial, and coronal MR Images (C,D). HPE given as osteosarcoma.




Figure 2: 45 yrs/F c/o swelling in the right hip since 5 months A: X-ray Pelvis with both hips shows Right iliac blade shows ill-defined lytic areas with a large soft tissue component with calcifications. B: CT axial section shows ill-defined lytic lesion with significant soft tissue component and multiple calcifications. C to D (Axial and Coronal STIR): Large multilobulated lesion noted




Figure 3: 17 yrs./F c/o pain in right ankle since 3 months A: XRAY -Right ankle (AP)- a well-defined subarticular lytic lesion with narrow zone of transition noted in the lower end of right tibia. There is no evidence of periosteal reaction – consistent with Osteoclastoma. B & C (T1 SAG & T2 SAG): A well-defined altered signal intensity lesion noted in the lower end of tibia which is hypointense to marrow on T1WI and hyperintense on T2WI extending into the subchondral region. D: A high power view reveals the giant cell nuclei and stromal cell nuclei to be similar, characteristic of GCT.




Figure 4: 14 yrs./M c/o pain and swelling in left thigh since 2 months A: XRAY -Left thigh (AP)- mild expansion with lytic and sclerotic areas with new bone formation and thick periosteal reaction with codman's triangle s/o osteogenic sarcoma. B & C (COR T2 FS & COR T1): Mildly widened with marrow showing iso to hypointense signal intensity on T1WI and heterogeneous (predominantly hyperintense) signal intensity on T2WI and STIR with areas of bone destruction showing spiculated periosteal reaction and new bone formation. D: H&E stained slide shows sheets of cells with atypia and mitoses with areas of atypical osteoid and few giant cells – s/o Osteosarcoma.



Figure 5: 15 yrs./F c/o pain and swelling in right thigh since 2 months. **A:** XRAY –Left thigh (AP)– expansile lesion with permeative type of bone destruction and onion peel like periosteal reaction with a soft tissue swelling. Features s/o Ewing's sarcoma. **B & D (COR T2WI & SAG T1WI):** Mild expansion with altered signal intensity in the marrow with lamellated periosteal reaction giving an onion peel appearance and a large lobulated soft tissue component associated with the lesion which shows heterogeneous signal intensity on both T1 and T2WI with areas of haemorrhage and necrosis. **C:** shows solid sheet of small round cells with indistinct cell borders, scant cytoplasm round to oval nucleus with vesicular chromatin with small nucleoli –s/o Ewing's sarcoma.

CONCLUSION

MRI in combination with plain radiography is an excellent modality for evaluation of the musculoskeletal pathologies especially differentiating a malignant from a benign lesion. The multiplanar imaging capabilities place a major role in delineation of tumour extent in to the bone and soft tissues with high contrast and resolution with additional information of neurovascular bundle involvement, joint involvement and staging.

REFERENCES

1. Weber K, Damron TA, Frassica FJ, Sim FH. Malignant bone tumors. Instr Course Lect. Departments of Orthopaedics and Oncology, Johns Hopkins University, Baltimore, Maryland, USA. 2008;57:673-88.
2. Morphology code of the International Classification of Diseases for Oncology (ICD-O) {726} and the Systematized Nomenclature of Medicine (<http://snomed.org>)
3. Von Eisenhart-Rothe R, Toepfer A, Salzmann M, Schauwecker J, Gollwitzer H, Rechl H. Primary malignant bone tumors; Orthopaedia. 2011 Dec;40(12):1121-42
4. Theodore T. Miller, MD. Bone Tumors and Tumorlike Conditions: Analysis with Conventional Radiography. From the Department of Radiology and Imaging, Hospital for Special Surgery, 535 E 70th St, New York, NY 10021; doi : 10.1148/ radiol. 2463061038 March 2008 Radiology, 246, 662-674.
5. Coleman RE, Holen I. Bone metastases. In: Abeloff MD, Armitage JO, Niederhuber JE, Kastan MB, McKenna WG, eds. Clinical Oncology. 4th ed. Philadelphia, Pa: Elsevier Churchill Livingstone; 2008:chap 57
6. Zeng Hui, Kang Bin, Liu Guoping ; Ultrasonographic diagnosis of bone tumors and its clinical implications. Tang Xingyu September 2001, Volume 21, Issue 3, pp 236-23.
7. Rydberg, J, Buckwalter KA, Caldemeyer KS, et al. Multisection CT: scanning techniques and clinical applications. Radiographics. 2000; 20:1787.
8. Colleen M. Costelloe, William A. Murphy Jr. and Beth A. Chasen Musculoskeletal Pitfalls in 18F-FDG PET/CT: Pictorial Review doi: 10.2214/AJR.07.7138 AJR September 2009 vol. 193 no. 3 Supplement WS1-WS13
9. Hailey D. Open magnetic resonance imaging (MRI) scanners. Issues Emerg Health Technol. 2006 Nov;(92):1-4.
10. Wintrobe MM, Lee GR. Wintrobe's Clinical Hematology. 10th ed. Baltimore, MD: Lippincott, Williams & Wilkins; 1999.
11. Enneking WF, Spawer SS, Goodman MA. A system for the surgical staging of musculoskeletal sarcoma. Clin Orthop 1980; 153:106.
12. Peterson H, Gillespie T, Hamlin DJ, Enneking WF, Springfield DS, Andrew ER, Spainer S, Slone R. Primary musculoskeletal tumours: Examination with MR imaging compared with conventional modalities. Radiology 1987;164:237-241.
13. Osolon PN, Everson L J , Griffiths H J . Staging of musculoskeletal tumours. RCNA 1999; 32:151-162.
14. Zimmer WD, Berquist TH, McLeod RA, Sim FH, Pritchard DJ, Shives TC, Word L E, May GR. Bone tumours : Magnetic Resonance versus Computer Tomography. Radiology 1985;155:709-718.
15. Shumann WP, Patten RM, Baron RL, Liddell RM, Conrad EV, Richardson ML. Comparison of STIR and Spin-Echo MR imaging at 1.5T in 45 suspected Extremity tumours : Lesion conspicuity and extent. Radiology 1991;179:247-252.
16. Erlmann R, Reiser MF, Peters PE, Vasallo P, Nommensen B, Kusnierz-Glanz CR, Ritter J, Roessner A. Musculoskeletal neoplasms : Static and Dynamic Gd-DTPA- enhanced MR imaging . Radiology 1989;171:767-773.
17. Norton KI, Hermann G, Abdelwahab IF, Klein MJ, Granowetter LF, Rabinowitz JG. Epiphyseal Involvement in Osteosarcoma. Radiology 1991;180:813-816.
18. Panicek DM, Gatsonis C, Rosenthal DI, Seegar LL, Huvo AG, Moore SG, Candry DJ, Palmer WE, McNeil BJ. CT and MR imaging in the local staging of Primary Malignant of Musculoskeletal Neoplasms: Report of Radiology Diagnostic Oncology group. Radiology 1997;202:739-744.
19. Loralic D Ma, Frassica FJ, Mc Carthy EF, Bhiemke DA, Zerhouni EA. Benign and Malignant Musculoskeletal masses : MR imaging Differentiation with Rim to center Differential Enhancement ratios. Radiology 1997;203:739-744.
20. Murali Sundaram, Mc Guire MH. Computed Tomography or Magnetic Resonance for evaluating the Solitary Tumour or Tumour like lesion of Bone? Skeletal Radiol 1988;17:393-401.