

A RANDOMIZED STUDY COMPARING PRE-OPERATIVE AND POST-OPERATIVE ENDOTHELIAL CELL COUNTS IN SINGLE-SITE VS. TWIN-SITE PHACOTRABECULECTOMY

Matta Namrata Kamal¹, Vijay Bhanudas Barge²

Received : 02/06/2026
Received in revised form : 01/07/2026
Accepted : 25/07/2026

Keywords:

Trabeculectomy, Glaucoma, Twin Site, IOP, Endothelial Cell Loss.

Corresponding Author:

Dr. Vijay Bhanudas Barge,
Email: bargevb@gmail.com

DOI: 10.47009/jamp.2026.8.4.105

Source of Support: Nil,
Conflict of Interest: None declared

Int J Acad Med Pharm
2026; 8 (4); 606-610



¹Consultant Ophthalmologist, Siddhigiri Hospital & Research Centre, Kanheri, Kolhapur, Maharashtra, India.

²Assistant Professor, Department of Ophthalmology, Prakash Institute of Medical Sciences and Research, Ishwarpur, Sangli, Maharashtra, India.

ABSTRACT

Background: Phacotrabeculectomy is a combined procedure in which both cataract extraction and glaucoma filtering surgery are performed in the same operative session. The one-site approach consists of both cataract extraction and trabeculectomy occurring through a single scleral incision. The two-site approach begins with cataract surgery, typically through a clear cornea incision, followed by a completely separate trabeculectomy performed superiorly upon completing the cataract procedure. This study looks at IOP changes, vision, loss of endothelial cells and BCVA between single-site and twin-site phacoemulsification -trabeculectomy surgeries. **Materials and Methods:** This prospective, randomized study was conducted between September 2014 to August 2015 with sample size of 80 subjects at National Institute of Ophthalmology, Pune. **Results:** The endothelial cell loss was significantly more in single site surgery when compared to twin site surgery, in both at 1 month and 3 months follow-up time points (P-value<0.001, unpaired t test). BCVA improved by 50.82% in Single Site and 62.97 % in Twin Site when compared to Pre-op Visual Acuity at 3 months post-op time point. Distribution of subjects improved to the desired direction on BCVA in both single site and twin site procedures at both 1 month and 3-month follow-up time points. **Conclusion:** Phacotrabeculectomy surgery should be considered in patients with required low target IOP, complex medical regimens and advanced glaucoma. In terms of corneal safety twin site surgery has an edge over single site surgery at a follow up period of 3 months.

INTRODUCTION

The techniques of combined phacoemulsification-trabeculectomy are under continual development as new suture-less incisions, filtration procedures, and antimetabolite use are studied. The results of numerous studies show that the combined procedure is an effective method of treatment of glaucoma and cataract.^[1] Phacotrabeculectomy is a combined procedure in which both cataract extraction and glaucoma filtering surgery are performed in the same operative session. The one-site approach consists of both cataract extraction and trabeculectomy occurring through a single scleral incision. The two-site approach begins with cataract surgery, typically through a clear cornea incision, followed by a completely separate trabeculectomy performed superiorly upon completing the cataract procedure. Since, there are a few studies on loss of endothelial cells between single-site and twin-site

phacoemulsification -trabeculectomy surgeries, we undertook this study, in which we have also evaluated the post-operative Best corrected visual acuity (BCVA) and Intra Ocular Pressure (IOP) at 1 month and 3 months follow-up periods, in addition.

MATERIALS AND METHODS

This prospective, randomized study was conducted between September 2014 to August 2015 with sample size of 80 subjects at National institute of Ophthalmology, Pune. Surgeries were performed by a single surgeon. Subsequent to the diagnosis of glaucoma and cataract the patient was counseled about the various treatment modalities available to him along with its benefits and complications. A formal consent was sought from the patient before the procedure was carried out under all aseptic precautions. The patient was then followed up at regular intervals for 6 months. The parameters

collected were Specular Microscopy, best corrected visual acuity (BCVA) and Intra Ocular Pressure. Inclusion criteria included patients from either gender, aged 55 to 85 years and Presence of visually significant cataract in patients with a. POAG which was not controlled by maximum drug therapy or patients with advanced disc damage; b. When there was appreciable visual field loss and optic disc damage with moderate IOP control (Mean Deviation ≥ 12); or c. Advanced field loss (Mean Deviation ≥ 18) where even transient IOP elevations might cause further optic nerve damage. Patients with endothelial pathology, preoperative inflammatory ocular disease, pseudo-exfoliation, angle closure disease / shallow anterior chamber, complicated cataract, more than grade III nuclear sclerosis, non – dilating pupil, previous intraocular surgery / trauma to same eye, and those who did not understand implication of protocol, and refused consent were excluded. Pre-operative examination consisted of BCVA, IOP measurement by Goldmann Applanation Tonometer, Slit-lamp examination, Dilated Fundus examination, and specular microscopy through non-contact Specular Microscope (Topcon). At follow-up visits i.e., at (Day 30 {1 Month} & Day 90 {3 Months}), Visual acuity, detailed slit-lamp examination for any post-op surgical complications, specular Microscopy for endothelial cell counts, and dilated Fundus examination. All readings were taken by single observers to obtain consistent results. For analysis, only ECC taken by non-contact Specular Microscope (Topcon) were taken into consideration. Procedure: Pre-operative preparations included written informed consent, Xylocaine sensitivity, and advised nil by mouth 2 hours before surgery. Dilatation of pupils was done with tropicamide (0.8%) and phenylephrine hydrochloride (0.5%) eye drops every 15 minutes for 4 times, 1 hour before surgery. Local Anesthesia with Plain Xylocaine 2% (peribulbar block) was done. Steps of Single site phacotrabeculectomy with mitomycin C (MMC) were, eye was painted and draped, eyelid was retracted with eye speculum, superior rectus bridal suture was taken, fornix based conjunctival flap was made, bleeders, if any were cauterized with electric cautery, triangular 4×4mm partial thickness scleral incision with limbal base was done with 15 number blade mounted on Bard- Parker handle. A weck-cel sponge with MMC (0.2mg/ml) was positioned on scleral flap for 2 min. After MMC application, area was thoroughly washed with 30 ml balanced salt solution. A limbal based triangular flap of two third of scleral thickness was dissected. Sclerocorneal tunnel was made by crescent. A keratome (2.8 mm) was used to enter the anterior chamber through sclerocorneal tunnel. Anterior chamber was formed with visco-elastic substance & continuous curvilinear capsulorrhexis was performed using a bent 26 no. needle. Phacoemulsification was performed using AMO Compact phacoemulsification machine. Stop and chop technique was used. The power setting was decided as per the grade of cataract. Cortical matter

was removed by bimanual irrigation and aspiration. A foldable lens was implanted in the bag. Side ports were sealed by hydration. The sclerostomy of 1.5 mm with Kelly's punch was made followed by peripheral iridectomy. Then scleral flap was repositioned and one fixed or releasable apical suture was taken with 10-0 nylon suture. Conjunctival flap closed with two watertight 8-0 vicryl sutures. Injection Moxifloxacin 0.5 mg in 0.1 cc given intra-camerally. Subconjunctival injection of Gentamycin & Dexamethasone was given inferonasally. Atropine ointment was applied & eye padded.

Steps of twin site phacotrabeculectomy with mitomycin C (MMC) were similar initial steps of single site procedure till a sclerocorneal tunnel by crescent. After that a separate superotemporal self-sealing clear corneal incision was taken (2.8 mm in size). Anterior chamber was formed with visco-elastic substance & continuous curvilinear capsulorrhexis was performed using a bent 26 no. needle. Phacoemulsification was done using AMO Compact phacoemulsification machine. Stop and chop technique was used. The power setting was decided as per the grade of cataract. Cortical matter was removed by bimanual irrigation and aspiration. A foldable lens was implanted in the bag. Wound was sealed by hydration. Now a keratome (2.8 mm) was used to enter the anterior chamber via previously dissected sclerocorneal tunnel. The sclerostomy of 1.5 mm with Kelly's punch was made followed by peripheral iridectomy. Then scleral flap was repositioned and one fixed or releasable apical 10-0 nylon suture was taken. Conjunctival flap closed with two water-tight 8-0 vicryl sutures. Injection Moxifloxacin 0.5 mg in 0.1 cc given intra-camerally. Subconjunctival injection of Gentamycin & Dexamethasone was given inferonasally. Atropine ointment was applied & eye padded. Post-operative follow up was on Day 30 (1 Month) and Day 90 (3 Months) respectively. Day

30 (1 month) and Day 90 (3 months) and follow up visit patients were assessed under following parameters: Thorough slit lamp examination for bleb morphology was performed at every visit & post-op. Operative bleb manipulation like argon laser suturolysis, releasable suture release and regular digital massage was performed accordingly. Best corrected visual acuity was recorded at each visit with Snellen's visual acuity chart. IOP was recorded with Goldmann Applanation Tonometer. Endothelial Cell Counts were done with Non-Contact Specular Microscope (Topcon) were assessed on Day 30 and day 90 follow-ups.

Statistical analysis: The data are presented using percentage s and for quantitative variables with mean and SD and range. The pre-op and post-op comparisons are done using Paired t test, after confirming the underlying normality assumption of difference of variables measured. A P value of less than 0.05 was set for statistical significance. Data was analyzed in SPSS (Statistical package for social sciences) version 19.0. The study protocol approved

by local ethical committee (National Institute of Ophthalmology) adhered to the tenets of the Declaration of Helsinki.

RESULTS

In the study 109 eyes of 80 patients were included out of which 59 eyes were operated by single site phacotrabeculectomy with MMC & 50 eyes were operated by twin site phacotrabeculectomy with MMC. [Table 1] shows the distribution of study subjects based on descriptive characteristics and other parameters among two study groups, single site and twin site. The average Log MAR visual acuity (Pre-op) and (1-month post-op follow-ups) differ significantly between study groups (P-value<0.001, unpaired t test). The endothelial cell loss was significantly more in single site surgery when compared to twin site surgery, in both at 1 month and 3 months follow-up time points (P-value<0.001, unpaired t test). BCVA improved by 50.82% in Single Site and 62.97 % in Twin Site when compared to Pre-op Visual Acuity at 3 months post-op time point. [Table 2] Distribution of subjects improved to the desired direction on BCVA in both single site and twin site procedures at both 1 month and 3-month follow-up time points. [Table 3]



Figure 1: Single Site surgery



Figure 2: Twin-site phacotrabeculectomy

Table 1: Distribution of study subjects based on descriptive characteristics and other parameters among single-site or twin-site

Variable		Single site		Twin site		P value
		n	%	n	%	
Age-group in years	55-64	14	23.7	7	14	
	65-74	27	45.8	29	58	
	75-84	18	30.5	14	28	
Gender	Male	32	54.23	39	78	0.015
	Female	27	45.77	11	22	
Laterality	Right eye	31	52.54	25	50	0.849
	Left eye	28	47.46	25	50	
Quantitative		Mean	SD	Mean	SD	
Age in years		70.07	7.59	70.74	6.16	0.611
Pre-op vision (Log Mar)		0.62	0.21	0.85	0.39	<0.001
Pre-op IOP mmHg		24.85	3	24.86	2.58	0.274
Pre-op ECC/mm ²		2230.61	243.91	2155.94	323.50	0.183

Table 2: Inter group Comparison of pre-operative and post-operative Vision, IOP, Endothelial Cell Counts and relative changes between two study groups at different time points

Variable	Follow-up	Single Site Group (n=59)	Twin Site Group (n=50)	P-value
Vision (Log MAR)	Pre-op	0.62 ± 0.21	0.85 ± 0.39	<0.001
	Post Op 1 month	0.50±0.18	0.60 ± 0.28	0.038
	Post op 3 Month	0.31±0.14	0.31 ± 0.17	0.769
IOP (mmHg)	Pre-op	24.85 ± 3.00	24.26 ± 2.58	0.274
	Post Op 1 month	13.15 ± 1.76	13.24 ± 2.47	0.835
	Post op 3 Month	11.73 ± 1.19	11.90 ± 1.05	0.427
ECC/mm ²	Pre-op	2230.61 ± 243.91	2155.94 ± 323.50	0.183
	Post Op 1 month	2079.46 ± 235.99	2056.88 ± 316.73	0.706
	Post op 3 Month	1980.75 ± 232.99	1978.94 ± 302.93	0.967

Endothelial cell loss	at 1 month	151.15 ± 58.17	99.06 ± 54.93	< 0.001
	at 3 months	249.86 ± 62.44	177.0 ± 114.99	< 0.001
Relative % change with respect to preop value in vision	at 1 month	19.67	29.72	
	at 3 months	50.82	62.97	
Relative % change with respect to preop value in IOP	at 1 month	47.07	45.42	
	at 3 months	52.80	50.95	
Relative % change with respect to preop value in Endothelial cell loss	at 1 month	6.78	4.59	
	at 3 months	11.2	8.21	

Table 3: Distribution of patients according to best corrected visual acuity (BCVA) single site and Twin site Phacotrabeculectomy

	Group	BCVA				Total
		≥ 6/9	6/12 - 6/18	6/24 - 6/36	≤ 6/60	
Pre-operative	Single site	0	25	26	8	59
	Twin site	0	13	17	20	50
At 1 month	Single site	2	46	9	2	59
	Twin site	1	29	12	8	50
At 3 months	Single site	21	35	1	1	58
	Twin site	19	29	0	2	50

DISCUSSION

Since the first phacoemulsification procedure performed by Kelman,^[2] in 1967, corneal endothelial cell loss (ECL) remains a serious concern in cataract surgery.^[3] Damage to the corneal endothelium is influenced by various pre- and intra-operative factors.^[3] In phacoemulsification, the amount of ECL determines final postoperative corneal transparency and to some extent visual acuity. Thus, attempts to protect the corneal endothelium and to minimize their damage can play an important role in improving quality of life in patients with cataracts.^[4-5] Risk of endothelial cell damage increases with a high nucleus grade, advanced age, long phacoemulsification time (Phaco time), high ultrasound energy, small pupil diameter, large infusion volume, type of intraocular lens (IOL), and short axial length.^[2-8] Hence, a surgeon who is more familiar with temporal entry for phacoemulsification, will be causing less endothelial cell damage in twin site surgery, the steps of trabeculectomy essentially being the same for both single site and twin site surgeries.

A study by Nassiri et al, which is a historical cohort study comparing corneal endothelial cell loss in single site and twin site phacotrabeculectomy. In both groups, the longer the follow-up time was, the higher the corneal endothelial cell loss and the corneal endothelial cell area expansion was. One-site surgery seems to cause less corneal endothelial cell damage than the 2-site operation during the median follow-up time of 1 year, while in our study the, mean corneal endothelial cell density (±SD) in single site group, at a follow up period of 3 months was 1980.75±232.99/mm², while that in twin site group was 1978.94±302.93/mm², with no statistically significant difference between the two.^[9]

The study by Soro Martinez et al., which was an observational retrospective study, compares phacoemulsification trabeculectomy + intraocular lens implantation with trabeculectomy alone. Here,

80 eyes of 62 patients were divided into one control group and three experimental groups, in which trabeculectomy or phacotrabeculectomy and IOL implantation were performed. Mean corneal endothelial cellular density (±SD) was 2,619 ±319/mm² in the control group, 2,447 ±425/mm² in the trabeculectomy group, 1,968 ±342/mm² in the one-step phacotrabeculectomy group, and 1,551 ±323/mm² in the two-step phacotrabeculectomy group. This study states that corneal endothelial cell loss is more in combined surgery and that two-step procedure causes more endothelial cell loss than single step surgery. In our study, mean corneal endothelial cellular density (±SD) was 2,230.61±243.91/mm² in single site and 2155.94±323.50/mm² in twin site phacotrabeculectomy, while the post-operative mean corneal endothelial cellular density at 3 months was 1980.75±232.99/mm² in single site and 1982.94±303.49/mm² in twin site phacotrabeculectomy. The endothelial cell loss (at a follow up period of 3 months) is 249.86±62.44/mm² in single site and 177.00±114.99/mm² in twin site surgery. Hence, there is a significantly lower loss of corneal endothelial cells in twin site phacotrabeculectomy as compared to single site. In the present study, the visual acuity after a follow up period of 3 months improved both in single site and twin site to values being 50.82% in single site and 62.97% in twin site surgeries and IOP was reduced to an average of 11.81±1.13 at a follow up period of 3 months.^[10]

A prospective randomized controlled trial in 2008 by Buys YM et al., noted no statistically significant difference in post-operative IOP between two types of surgeries. Corneal endothelial cell counts were significantly lower in the twin-site group at 3 and 12 months.^[11] Another study suggested that patients with glaucoma may have lower corneal endothelial cell density compared to those without glaucoma of the same age group. The proposed mechanisms are direct damage from IOP, congenital alteration of the

corneal endothelium in patients with glaucoma, glaucoma medication toxicity, or a combination of these.^[12]

A single-site surgery is likely faster to perform than a twin-site procedure, but more recent studies have begun to show that not only does a twin-site procedure produce better post-operative IOP control but also produce less need for adjunct anti glaucoma medications. A twin-site procedure also has safety advantages over a single-site technique as there is less risk of anti-metabolites entering the anterior chamber, and less manipulation of the conjunctiva and sclerostomy. Moreover, most of the surgeons are familiar with temporal entry for routine phacoemulsifications, hence they invariably cause lesser damage to the endothelium in case of twin site surgeries.

CONCLUSION

Phacotrabeculectomy surgery should be considered in patients with required low target IOP, complex medical regimens and advanced glaucoma. In terms of corneal safety, twin site surgery has an edge over single site surgery at a follow up period of 3 months. The results of twin site and single site phacotrabeculectomy in terms of IOP and visual outcome are comparable at a follow up period of 3 months.

REFERENCES

1. Hsu CH, Obstbaum SA. Technique and outcome of combined phacoemulsification and trabeculectomy. *Curr Opin Ophthalmol*. 1998;9(2):9-14.
2. Kelman CD. Phacoemulsification and aspiration; A new technique of cataract removal: a preliminary report. *Am J Ophthalmol*. 1967;64:23-35.
3. Walkow T, Anders N, Klebe S. Endothelial cell loss after phacoemulsification: relation to preoperative and intraoperative parameters. *J Cataract Refract Surg*. 2000;26:727-732.
4. Díaz-Valle D, Benítez del Castillo Sánchez JM, Castillo A, Sayagués O, Moriche M. Endothelial damage with cataract surgery techniques. *J Cataract Refract Surg*. 1998;24:951-955.
5. Dick HB, Kohnen T, Jacobi FK, Jacobi KW. Long-term endothelial cell loss following phacoemulsification through a temporal clear corneal incision. *J Cataract Refract Surg*. 1996;22:63-71.
6. Dosso AA, Cottet L, Burgener D, Di Nardo S. Outcomes of coaxial microincision cataract surgery versus conventional coaxial cataract surgery. *J Cataract Refract Surg*. 2008;34:284-288.
7. Storr-Paulsen A, Norregaard JC, Ahmed S, et al. Endothelial cell damage after cataract surgery: divide-and conquer versus phaco-chop technique. *J Cataract Refract Surg*. 2008;34:996-1000.
8. Walkow T, Anders N, Klebe S. Endothelial cell loss after phacoemulsification: relation to preoperative and intraoperative parameters. *J Cataract Refract Surg*. 2000;26:727-732.
9. Nassiri N, Nassiri N, Majidi-N M, et al. Corneal endothelial cell changes after Ahmed valve and Molteno glaucoma implants. *Ophthalmic Surgery, Lasers and Imaging Retina* 2011;42(5):394-399.
10. Soro-Martínez MI, Villegas-Pérez MP, Sobrado-Calvo P, Ruiz-Gómez JM, Mora-Figueroa JI. Corneal endothelial cell loss after trabeculectomy or after phacoemulsification, IOL implantation and trabeculectomy in 1 or 2 steps. *Graefes Arch Clin Exp Ophthalmol*. 2010;248(2):249-56.
11. Buys YM, Chipman ML, Zack B, Rootman DS, Lomovic AR, Trope GE. Prospective randomized comparison of one- versus two-site Phacotrabeculectomy two-year results. *Ophthalmology*. 2008;115(7):1130-1133.e1.
12. Gagnon MM, Boisjoly HM, Brunette I, Charest M, Amyot M. Corneal endothelial cell density in glaucoma. *Cornea*. 1997;16(3):314-8.